

Five Things Physicians and Patients Should Question

1 **Don't obtain baseline laboratory studies in patients without significant systemic disease (ASA I or II) undergoing low-risk surgery – specifically complete blood count, basic or comprehensive metabolic panel, coagulation studies when blood loss (or fluid shifts) is/are expected to be minimal.**

Performing routine laboratory tests in patients who are otherwise healthy is of little value in detecting disease. Evidence suggests that a targeted history and physical exam should determine whether pre-procedure laboratory studies should be obtained. The current recommendation from the 2003 ASA amendment that all female patients of childbearing age be offered pregnancy testing rather than required to undergo testing has provided individual physicians and hospitals the opportunity to set their own practices and policies relating to preoperative pregnancy testing. Some institutions respect the right of a patient to refuse testing after a thorough explanation of the anesthetic risks during pregnancy and the required signing of a waiver. The avoidance of the routine administration of the pregnancy test was therefore excluded from our Top 5 preoperative recommendations.

The risk specifically related to the surgical procedure could however modify the above preoperative recommendation to obtain laboratory studies and when the need arises; the decision to implement should include a joint decision between the anesthesiologists and surgeons. This should be applicable to all outpatient surgery.

2 **Don't obtain baseline diagnostic cardiac testing (trans-thoracic/ esophageal echocardiography – TTE/TEE) or cardiac stress testing in asymptomatic stable patients with known cardiac disease (e.g., CAD, valvular disease) undergoing low or moderate risk non-cardiac surgery.**

Advances in cardiovascular medical management, particularly the introduction of perioperative beta-blockade and improvements in surgical and anesthetic techniques, have significantly decreased operative morbidity and mortality rates in noncardiac surgery. Surgical outcomes continue to improve causing the mortality rate of major surgeries to be low and the need for revascularization minimal. Consequently, the role of preoperative cardiac stress testing has been reduced to the identification of extremely high-risk patients, for instance, those with significant left main disease for which preoperative revascularization would be beneficial regardless of the impending procedure. In other words, testing may be appropriate if the results would change management prior to surgery, could change the decision of the patient to undergo surgery, or change the type of procedure that the surgeon will perform.

3 **Don't use pulmonary artery catheters (PACs) routinely for cardiac surgery in patients with a low risk of hemodynamic complications (especially with the concomitant use of alternative diagnostic tools (e.g., TEE).**

The increased risk of hemodynamic complications as indicated above is defined as a patient with clinical evidence of significant cardiovascular disease; pulmonary dysfunction, hypoxia, renal insufficiency or other conditions associated with hemodynamic instability (e.g., advanced age, endocrine disorders, sepsis, trauma, burns).

The use of a PAC during cardiac surgery has been associated with increased mortality and a higher risk of severe end-organ complications. There is clear consensus in the literature that the use of a PAC cannot be recommended as a matter of routine, but for a definite role in a very select group of patients undergoing cardiac surgery. According to a survey by practicing anesthesiologists, the use of PAC could be recommended for specific indications in cardiac surgery including coronary artery bypass grafting (CABG) with poor left ventricular (LV) function, LV aneurysmectomy, recent myocardial infarction, pulmonary hypertension, diastolic dysfunction, acute ventricular septal rupture and insertion of left ventricular assist device. The appropriate indications remain debatable. However, although the PAC has no role in routine perioperative care, the existence of a specific subpopulation for which the use of this device may be beneficial cannot be excluded.

4

Don't administer packed red blood cells (PRBCs) in a young healthy patient without ongoing blood loss and hemoglobin of ≥ 6 g/dL unless symptomatic or hemodynamically unstable.

The hemoglobin transfusion threshold used in multiple studies has varied from 6.0 to 10.0 g/dL. The optimal hemoglobin/hematocrit criterion for transfusion remains controversial in several clinical settings. Nevertheless, compared with higher hemoglobin thresholds, a lower hemoglobin threshold is associated with fewer red blood cell units transfused without adverse associations with mortality, cardiac morbidity, functional recovery or length of hospital stay. Hospital mortality remains lower in patients randomized to a lower hemoglobin threshold for transfusion versus those randomized to a higher hemoglobin threshold.

The decision to transfuse should be based on a combination of both clinical and hemodynamic parameters.

5

Don't routinely administer colloid (dextrans, hydroxyethyl starches, albumin) for volume resuscitation without appropriate indications.

There is no evidence from multiple randomized controlled trials and recent reviews/meta-analyses that resuscitation with colloids reduces the risk of death compared to crystalloids. Colloids offer no survival benefit and are considerably more expensive than crystalloids; their continued routine use in clinical practice should therefore be questioned. Recent perioperative data on the use of colloids in certain populations remain controversial; nevertheless, there is consensus on the avoidance of the routine use of colloids for volume resuscitation in the general surgical population given the overwhelming amount of evidence in the literature of possible harm when used in un-indicated patients. Health care providers should refer to the current evolving literature when faced with specific conditions like sepsis, traumatic brain injury, acute renal injury and burns thereby creating a forum for discussion among the care providers of the efficacy of such a treatment in that individual patient.

Nevertheless, it is important to note that the endpoint in most studies is mortality and morbidity. There is insufficient data to adequately address the need of colloids over crystalloids for other endpoints of interest like hypotension, need for blood transfusion, length of hospital stay, etc. Further research may be required to delineate the existence of any particular benefits of colloids over crystalloids.

How This List Was Created

The list started as an academic project of Onyi C. Onuoha, M.D., M.P.H. A review of the literature and practice guidelines as approved by the American Society of Anesthesiologists (ASA) was performed to identify an evidence-based list of activities to question within the field of anesthesiology. A multi-step survey of anesthesiologists in both the academic and private sector and ASA Committees of Jurisdiction was performed to generate a “Top 5 List” list of preoperative and intraoperative activities. The final list was endorsed by the ASA and accepted for the *Choosing Wisely*[®] campaign. We believe that developing strategies whereby all stakeholders in the perioperative team are involved in the implementation is a means in which anesthesiologists could be engaged in the efforts to reduce over-utilization of low value, non-indicated medical services evident in the U.S. health system today.

ASA's disclosure and conflict of interest policy can be found at www.asahq.org.

Sources

Committee on Standards and Practice Parameters, Apfelbaum JL, Connis RT, Nickinovich DG; American Society of Anesthesiologists Task Force on Preanesthesia Evaluation, Pasternak LR, Arens JF, Caplan RA, Connis RT, Fleisher LA, Flowerdew R, Gold BS, Mayhew JF, Nickinovich DG, Rice LJ, Roizen MF, Twersky RS. Practice advisory for preanesthesia evaluation: an updated report by the American Society of Anesthesiologists Task Force on Preanesthesia Evaluation. *Anesthesiology*. 2012 Mar;116(3):522–38.

Kumar A, Srivastava U. Role of routine laboratory investigations in preoperative evaluation. *J Anaesthesiol Clin Pharmacol*. 2011;27(2):174–9.

Mollov JL, Twersky RS. (2013). Is routine preoperative pregnancy testing necessary? In L. Fleisher, Evidence-based practice of anesthesiology (3rd ed., pp. 26-30). Philadelphia (PA): Elsevier Saunders.

Soares Dde S, Brandao RR, Mourao MR, Azevedo VL, Figueiredo AV, Trindade ES. Relevance of routine testing in low risk patients undergoing minor and medium surgical procedures. *Rev Bras Anesthesiol*. 2013;63(2):197–201.

Brown SR, Brown J. Why do physicians order preoperative test? A qualitative study. *Fam Med*. 2011;43(5):338–43.

Czoski-Murray C, Lloyd JM, McCabe C, Claxton K, Oluboyede Y, Roberts J, Nicholls JP, Rees A, Reilly CS, Young D, Fleming T. What is the value of routinely testing full blood count, electrolytes and urea, and pulmonary function test before elective surgery in patients with no apparent clinical indication and in subgroups of patients with common comorbidities: a systematic review of the clinical and cost-effective literature. *Health Technol Assess*. 2012;16(50):1–159.

Katz RI, Dexter F, Rosenfeld K, Wolfe L, Redmond V, Agarwal D, Salik I, Goldstein K, Goodman M, Glass PS. Survey study of anesthesiologists' and surgeons' ordering of unnecessary preoperative laboratory tests. *Anesth Analg*. 2011;112(1):207–12.

Keay L, Lindsley K, Tielsch J, Katz J, Schein O. Routine preoperative testing for cataract surgery. *Cochrane Database Syst Rev*. 2012;3:CD007293.

Committee on Standards and Practice Parameters, Apfelbaum JL, Connis RT, Nickinovich DG; American Society of Anesthesiologists Task Force on Preanesthesia Evaluation, Pasternak LR, Arens JF, Caplan RA, Connis RT, Fleisher LA, Flowerdew R, Gold BS, Mayhew JF, Nickinovich DG, Rice LJ, Roizen MF, Twersky RS. Practice advisory for preanesthesia evaluation: an updated report by the American Society of Anesthesiologists Task Force on Preanesthesia Evaluation. *Anesthesiology*. 2012 Mar;116(3):522–38.

Miller AL, Beckman JA. (2013). Which patient should have a preoperative cardiac evaluation (stress test)? In L. Fleisher, Evidence-based practice of anesthesiology (3rd ed., pp. 61–70). Philadelphia (PA): Elsevier Saunders.

Schiefermueller J, Myerson S, Handa AI. Preoperative assessment and perioperative management of cardiovascular risk. *Angiology*. 2013;64(2):146–50.

Sheffield KM, McAdams PS, Benarroch-Gampel J, Goodwin JS, Boyd CA, Zhang D, Riall TS. Overuse of preoperative cardiac stress testing in medicare patients undergoing elective noncardiac surgery. *Ann Surg*. 2013; 257(1):73–80.

Almanaseer Y, Mukherjee D, Kline-Rogers EM, Kesterson SK, Sonnad SS, Roges B, Smith D, Furney S, Ernst R, McCort J, Eagle KA. Implementation of the ACC/AHA guidelines for preoperative cardiac risk assessment in a general medicine preoperative clinic: improving efficiency and preserving outcomes. *Cardiology*. 2005;103(1):24–9.

Cinello M, Nucifora G, Bertolissi M, Badano LP, Fresco C, Gonano N, Fioretti PM. American College of Cardiology/American Heart Association perioperative assessment guidelines for noncardiac surgery reduces cardiologic resource utilization preserving favorable outcome. *J Cardiovasc Med*. 2007;8(11):882–8.

Augoustides JG, Neuman MD, Al-Ghofaily L, Silvay G. Preoperative cardiac risk assessment for noncardiac surgery: defining costs and risks. *J Cardiothorac Vasc Anesth*. 2013;27(2):395–9.

Falcone RA, Nass C, Jermyn R, Hale CM, Stierer T, Jones CE, Walters GK, Fleisher LA. The value of preoperative pharmacologic stress testing before vascular surgery using ACC/AHA guidelines: a prospective randomized trial. *J Cardiothorac Vasc Anesth*. 2003;17(6):694–8.

Poldermans D, Boersma E. Beta-blocker therapy in noncardiac surgery. *N Engl J Med*. 2005;353:412–4.

American Society of Anesthesiologists Task Force on Pulmonary Artery Catheterization. Practice guidelines for pulmonary artery catheterization. *Anesthesiology*. 2003 Oct; 99:988–1014.

Schwann NM, Hillel Z, Hoelt A, Barash P, Mohnle P, Miao Y, Mangano DT. Lack of effectiveness of the pulmonary artery catheter in cardiac surgery. *Anesth Analg*. 2011;113(5):994–1002.

Rajaram SS, Desai NK, Kalra A, Gajera M, Cavanaugh SK, Brampton W, Young D, Harvey S, Bowan K. Pulmonary artery catheters for adult patients in intensive care. *Cochrane Database Syst Rev*. 2013;2:CD003408.

Kanchi M. Do we need a pulmonary artery catheter in cardiac anesthesia? – An Indian perspective. *Ann Card Anaesth*. 2011;14(1):25–9.

Harvey S, Stevens K, Harrison D, Young D, Brampton W, McCabe C, Singer M, Rowan K. An evaluation of the clinical and cost-effectiveness of pulmonary artery catheters in patient management in intensive care: a systematic review and a randomized controlled trial. *Health Technol Assess*. 2006;10(29):1–133.

Ramsey SD, Saint S, Sullivan SD, Day L, Kelley K, Bowdye A. Clinical and economic effects of pulmonary artery catheterization in nonemergent coronary artery bypass surgery. *J Cardiothorac Vasc Anesth*. 2000;14(2):113–8.

Chatterjee K. Historical Perspectives in Cardiology. The Swan-Ganz catheters: past, present, and future – a viewpoint. *Circulation*. 2009;119:147–52.

Sandham JD, Hull RD, Brant RF, Knox L, Pineo GF, Doig CJ, Laporta DP, Viner S, Passerini L, Devitt H, Kirby A, Jacka M; Canadian Critical Care Clinical Trials Group. A randomized, controlled trial of the use of pulmonary-artery catheters in high-risk surgical patients. *N Engl J Med*. 2003;348:5–14.

Miller AL, Beckman JA. (2013). Which patient should have a preoperative cardiac evaluation (stress test)? In L. Fleisher, Evidence-based practice of anesthesiology (3rd ed., pp. 61–70). Philadelphia (PA): Elsevier Saunders.

American Society of Anesthesiologists Task Force on Perioperative Blood Transfusion and Adjuvant Therapies. Practice guidelines for perioperative blood transfusion and adjuvant therapies. *Anesthesiology*. 2006 Jul;105(1):198–208.

Carson JL, Carless PA, Hebert PC. Outcomes using lower versus higher hemoglobin thresholds for red blood cell transfusion. *JAMA*. 2013;309(1):83–4.

Carson JL, Patel MS. (2013). Is there an optimal perioperative hemoglobin level? In L. Fleisher, Evidence-based practice of anesthesiology (3rd ed., pp. 155–163). Philadelphia (PA): Elsevier Saunders.

Goodnough LT, Levy JH, Murphy MF. Concepts of blood transfusion in adults. *Lancet*. 2013;381(9880):1845–54.

Carson JL, Carless PA, Hebert PC. Transfusion threshold and other strategies for guiding allogeneic red blood cell transfusion. *Cochrane Database Syst Rev*. 2012; 4:CD002042.

Bittencourt R, Costa J, Lobo JE, Aquiar FC. Consciously transfusion of blood products. Systematic review of indicative factors for blood components infusion trigger. *Rev Bras Anesthesiol*. 2012;62(3):402–10.

Carson JL, Grossman BJ, Kleinman S, Tinmouth AT, Marques MB, Fung MK, Holcomb JB, Illoh O, Kaplan LJ, Katz LM, Rao SV, Roback JD, Shander A, Tobian AA, Weinstein R, Swinton-McLaughlin LG, Djulbegovic B, Clinical Transfusion Medicine Committee of the AABB. Red blood cell transfusion: a clinical perspective guideline from the AABB. *Ann Intern Med*. 2012;157(1):49–58.

Toy P, Feiner J, Viele MK, Watson J, Yeap H, Weiskopf RB. Fatigue during acute isovolemic anemia in healthy resting humans. *Transfusion*. 2000;40(4):457–60.

Committee on Standards and Practice Parameters, Apfelbaum JL, Connis RT, Nickinovich DG; American Society of Anesthesiologists Task Force on Preanesthesia Evaluation, Pasternak LR, Arens JF, Caplan RA, Connis RT, Fleisher LA, Flowerdew R, Gold BS, Mayhew JF, Nickinovich DG, Rice LJ, Roizen MF, Twersky RS. Practice advisory for preanesthesia evaluation: an updated report by the American Society of Anesthesiologists Task Force on Preanesthesia Evaluation. *Anesthesiology*. 2012 Mar;116(3):522–38.

Perel P, Roberts I, Pearson M. Colloid versus crystalloid for fluid resuscitation in critically ill patients (Review). The Cochrane Collaboration, the Cochrane Library 2009;3.

Perel P, Roberts I, Ker K. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev*. 2013 Feb 28;2.

Perel P, Roberts I. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev*. 2012 Jun 13;6.

Perel P, Roberts I. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev*. 2011 Mar 16;(3):CD000567.

Perel P, Roberts I. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev*. 2007 Oct 17;(4):CD000567.

Roberts I, Alderson P, Bunn F, Chinnock P, Ker K, Schierhout G. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev*. 2004 Oct 18;(3):CD000567.

Kruer RM, Ensor CR. Colloids in the intensive care unit. *Am J Health Syst Pharm*. 2012 Oct 1;69(19):1635–42.

NATA: Network for Advancement and Transfusion Alternatives. Crystalloids versus colloids: the controversy [Internet]. NATA. 2013 [cited 2013 Sep 20]. Available from: <http://www.nataonline.com/np/158/crystalloids-versus-colloids-controversy>.

Reinhart K, Perner A, Sprung CL, Jaeschke R, Schortgen F, Johan Groeneveld AB, Beale R, Hartog CS; European Society of Intensive Care Medicine. Consensus statement of the ESICM task force on colloid volume therapy in critically ill patients. *Intensive Care Med*. 2012;38(3):368–83.

About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.

To learn more about the ABIM Foundation, visit www.abimfoundation.org.



About the American Society of Anesthesiologists

The American Society of Anesthesiologists (ASA) is an educational research and scientific association of physicians organized to raise and maintain the standards of the medical practice of anesthesiology and improves the care of the patient. Since its founding in 1905, the Society's achievements have made it an important voice in American medicine and the foremost advocate for all patients who require anesthesia or relief from pain. As physicians, anesthesiologists are responsible for administering anesthesia to relieve pain and for managing vital life functions, including breathing, heart rhythm and blood pressure, during surgery. After surgery, they maintain the patient in a comfortable state during the recovery and are involved in the provision of critical care medicine in the intensive care unit.

For more information about ASA, visit www.asahq.org.

